

Bibliografia

Bernstein A. et al., Major cereal grains fibers and psyllium in relation to cardiovascular health, *Nutrients* 2013; 5:1471.

Broekaert WF et al., Prebiotic effects of cereal-derived arabinoxylans, arabinoxylan-oligosaccharides and xylo-oligosaccharides, *Crit Rev Food Sci Nutr* 2011; 51: 178.

Boll, E. V. J., Ekström, L. M., Courtin, C. M., Delcour, J. A., Nilsson, A. C., Björck, I. M., & Östman, E. M. (2015). Effects of wheat bran extract rich in arabinoxylan oligosaccharides and resistant starch on overnight glucose tolerance and markers of gut fermentation in healthy young adults. *European journal of nutrition*, 1-10.

Cloetens L. et al., Tolerance of arabinoxylan- oligosaccharides and their prebiotic activity, *Brit J Nutr* 2010; 103-703

Coelho E et al., Revisiting the structural features of arabinoxylans from brewer's spent grain, *Carbohydrate Polymers* 2016; 139:167.

Damen B et al., Consumption of breads containing in Situ-Arabinoxylan Oligosaccharides Alters Gastrointestinal Effects in Healthy Volunteers, *J. Nutr.* 2012; 142:470.

Francois I et al. Effects of a wheat brain extract containing arabinoxylan oligosaccharides (WBE) on gastrointestinal health parameters in healthy adult human volunteers: a double-blind, randomised, placebo-controlled, cross over trial, *British Journal of Nutrition* 2012; 108:2229

Garcia, A. L., Steiniger, J., Reich, S. C., Weickert, M. O., Harsch, I., Machowetz, A., ... & Doerfer, J. (2006). Arabinoxylan fibre consumption improved glucose metabolism, but did not affect serum adipokines in subjects with impaired glucose tolerance. *Hormone and metabolic research*, 38(11), 761-766.

Garcia, A. L., Otto, B., Reich, S. C., Weickert, M. O., Steiniger, J., Machowetz, A., ... & Meuser, F. (2007). Arabinoxylan consumption decreases postprandial serum glucose, serum insulin and plasma total ghrelin response in subjects with impaired glucose tolerance. *European Journal of Clinical Nutrition*, 61(3), 334-341.

Giulia Falchi, A., Grecchi, I., Muggia, C., Palladini, G., & Perlini, S. (2016). Effects of a Bioavailable Arabinoxylan-enriched White Bread Flour on Postprandial Glucose Response in Normoglycaemic Subjects. *Journal of dietary supplements*, 13(6), 626-633.

Hartvigsen, M. L., Gregersen, S., Lærke, H. N., Holst, J. J., Knudsen, K. B., & Hermansen, K. (2014). Effects of concentrated arabinoxylan and β -glucan compared with refined wheat and whole grain rye on glucose and appetite in subjects with the metabolic syndrome: a randomized study. *European journal of clinical nutrition*, 68(1), 84-90.

Bibliografia

Hopkins M et al., Degradation of cross-linked arabinoxylans by the intestinal microbiota in children, *Appl Environ Microbiol* 2003; 69:6354

Lafiandra D et al. Improving cereal grain carbohydrates for diet and health, *J cereal Sci.* 2014; 59:312

Lu ZX, Walker KZ, Muir JG, Mascara T, O'Dea K. Arabinoxylan fiber, a byproduct of wheat flour processing, reduces the postprandial glucose response in normoglycaemic subjects. *Am J Clin Nutr.* 2000;71(5):1123–1128

McCarthy AL et al., Brewer's spent grain bioactivity of phenolic component its role on animal nutrition and potential for incorporation in functional foods: a review, *Proc Nutr Soc.* 2013; 72:117.

Malunga LN et al., Inhibition of alfa-glucosidase and glucose adsorption by feruloylated arabinoxylan mono and oligosaccharides from corn and wheat, *J. Nutr Metab* 2016; ID 1932532

Malunga LN et al., Isolation and Identification of feruloylated AX mono and oligosaccharides from undigested and digested maize and wheat, *Helyton* 2016; 2:e00106.

Malunga LN et al., Antihyperglycemic effect of water extractable AX from wheat aleurone and bran. *J Nutr* 2017; ID 5784759

Maki KC et al., Digestive and physiologic effects of a wheat bran extract, arabino-xylan- oligosaccharide in breakfast cereal, *Nutrition* 2012; 28:1115

Mendis M Et al. Arabinoxylans, gut microbiota and immunity, *Carbohydrate Polymers* 2016; 139:159

Mussatto L. Et al., Brewer's spent grain: a valuable feed stock for industrial application, *J SciFood Agric* 2014; 94:1264

Napolitano A et al., Potential prebiotic activity of oligosaccharides obtained by enzymatic conversion of durum wheat insoluble dietary fibre into soluble dietary fibre, *Nutr Metab Cardiovasc Dis* 2009; 19:283

Neyrink et al., Potential interest of gut microbial changes induced by non-digestible carbohydrates of wheat in the management of obesity and related disorders, *Curr Opin Clin Nutr Metab Care* 2010; 13:722.

Pollet A et al., In vitro fermentation of arabinoxylans from wheat bran and psyllium seed husk, *J Agric Food Chem* 2012; 60:946

Bibliografia

Reis S et al., Evaluation of the prebiotic potential of arabinoxylans from brewer's spent grain, Appl Microbiol Biotechnol 2014; 98:9365

Walton G et al., Gastrointestinal effects of arabinoxylan-enriched bread in healthy volunteers, Nutr J 2012; 11:36.

Zhang et al., Cereal-derived arabinoxylans as biological response modifiers: extraction, molecular features, and immune-stimulating properties, Crit Rev Food Sci Nutr. 2015; 55:1035.